

Case report

doi: <http://dx.doi.org/10.22265/acnef.0.0.310>

A true dialytic urgency: lithium intoxication

Intoxicación por litio, una verdadera urgencia dialítica

④ Mariana Montoya-Castillo^{1*}, ④ Tomás Giraldo-Hinestrosa², ④ Felipe Osorio Ospina³,
④ Diana Carolina Echeverri Diaz⁴, ④ María Antonia Rico-Mesa², ④ Carlos Aníbal Restrepo⁵,
④ Juan Simon Rico-Mesa⁶

¹ CES University, Medicine department, Medellín, Colombia

² CES University, School of Medicine, Medellín, Colombia

³ Pontifical Bolivarian University of Colombia, Medicine department, Medellín, Colombia

⁴ Cooperative University of Colombia, Medicine department, Medellín, Colombia

⁵ CES University, School of Medicine, Medellín, Colombia

⁶ Mayo Clinic, Rochester, Cardiology department, United States of America

Abstract

Lithium carbonate is a drug widely used in Colombia for the treatment of various psychiatric pathologies such as treatment in patients with bipolar affective disorder, however its narrow therapeutic index predisposes to the occurrence of acute poisoning, unlikely to happen but could be fatal.

It should be said that its diagnosis requires a high rate of suspicion because it presents various clinical manifestations such as neuropsychiatric symptoms and even mental disorders of an organic type that can be complicated by an acute kidney injury requiring emergency dialysis therapy.

We report the case of a patient who presented to the emergency department with alteration in the state of consciousness in whom lithium poisoning was found and who required hemodialysis in an early stage.

In conclusion, in a patient with risk factor and therapy with this drug is very important to guide appropriate treatment, and also discard differential diagnosis, in order to avoid fatal consequences.

Keywords: Dialysis, poisoning, lithium, chronic kidney disease, fatal outcome, urinary bladder diseases.

doi: <http://dx.doi.org/10.22265/acnef.0.0.310>

Resumen

El carbonato de litio es un fármaco ampliamente usado en Colombia para el tratamiento de varias patologías psiquiátricas, como es el caso del trastorno afectivo bipolar; sin embargo, su índice terapéutico estrecho predispone a intoxicación aguda. Aunque esta reacción rara vez ocurre, puede ser fatal.

El diagnóstico de intoxicación aguda por carbonato de litio requiere una alta tasa de sospecha, pues se presenta con varias manifestaciones como síntomas neuropsiquiátricos e incluso desórdenes mentales, que pueden complicarse por lesión renal aguda requiriendo terapia dialítica emergente.

A continuación se presenta el caso de una paciente que se llegó al servicio de urgencias con alteración en el estado de conciencia. Finalmente, se diagnosticó intoxicación por litio y la mujer requirió hemodiálisis.

En conclusión, en un paciente con factores de riesgo y terapia con este medicamento es muy importante guiar un manejo apropiado y descartar diagnósticos diferenciales, esto con el fin de evitar consecuencias fatales.

Palabras clave: diálisis, intoxicación; litio, enfermedad renal crónica, resultado fatal, enfermedades de la vejiga urinaria.

doi: <http://dx.doi.org/10.22265/acnef.0.0.310>



Citation: Montoya-Castillo M, Giraldo-Hinestrosa T, Osorio-Ospina F, Echeverri-Diaz DC, Rico-Mesa MA, Restrepo CA, Rico-Mesa JS. A true dialytic urgency: lithium intoxication. Rev. Colomb. Nefrol. 2018;5(2):179-189. doi: <http://dx.doi.org/10.22265/acnef.0.0.310>

*Correspondence: Mariana Montoya Castillo, marimontoya13@gmail.com

Received: 12-06-18 • Accepted: 26.07.18 • Published online: 9.08.18

Introduction

Lithium is a chemical element located in group one (alkali metal) of the periodic table. This specific location enables it to ionize into a cation and easily interact with other functional groups. In addition, it is the lightest metal (74 Daltons), making it easy to be distributed throughout the body¹. Its use in the treatment of mental disorders began in 1948 by Doctor John Cade in Australia². Since then, lithium salts were classified as mood stabilizers due to its positive empiric impact on the treatment of bipolar affective disorders with a success rate of 80% in the treatment of both maniac and depressive symptoms^{3,4}.

Case report

A 77 year old woman, with personal history of affective bipolar disorder in treatment with

lithium, presented into our hospital with functional declination, somnolence, fever and tremor, for 5 days approximately. The physical examination showed altered consciousness state, normal blood pressure, tachycardia, tachypnea, grade II dehydration, neck rigidity, tremor and hyperreflexia with clonus. The rest was normal.

Because the hospital was of first level, basic laboratories were available, so initially there was focus like meningitis, so antibiotics are started after blood and urine samples were collected. We observed leukocytosis with neutrophilia, altered renal function with GFR in 27.08 ml/min/1.73 m² with MDRD formula (**figure 1**), proteinuria, hematuria and massive bacteriuria in the urine sediment. Also glycemic levels and an electrocardiogram were realized, finding in the last one sinus tachycardia, normal ventricular axis, interval PR of 180 milliseconds, but a corrected interval QT of 510 milliseconds.




Figure 1. Laboratories

In this moment, with these findings, acute lithium intoxication was suspected associated with acute kidney injury and urinary tract infection, considered like an emergency dialysis. Immediately the patient was translated to a hospital with the service of nephrology and dialysis, and realized serum lithium levels finding an abnormal value of 2.85 mmol/L (Figure 1) (normal value 0.6 – 1.2 mmol/L). So,

it was decided to initiate dialysis femoral via and antibiotic, presenting good response. In urine culture was isolated *Proteus Mirabilis*, so regimen antibiotic with Aztreonam was gave during 10 days with resolution of the infection.

After three days of treatment, renal function and serum lithium level were normal.

Discussion

Lithium distribution volume ranges between 0.6-0.9 L/Kg in the average adult. It has no protein or tissue binding, making it a substance with a therapeutic index that ranges between 0.6-1.5 mEq/L, which is very narrow³. Particularly, the optimal steady-state concentration of lithium for treatment of bipolar disorders is 0.6-1.2 mEq/L. Lithium acts with inositol and glycogen synthase kinase 3, by inhibiting an enzyme called inositol monophosphatase, decreasing inositol levels and altering first and second messenger system. With the chronic administration, stabilize GABA reuptake, explaining the effect on mania. Also antagonize 5HT1A and 5HT1B receptors, increasing levels of serotonin, explaining the effect on depression symptoms. Maintaining these exact ranges may be challenging and chances of toxicity remain high. Lithium is excreted in its vast majority by the kidneys and because of its chemical resemblance to sodium; they are handled in a very similar way. It is freely filtered and 60% of it is reabsorbed in the proximal tubule⁵. Due to its high rate of reabsorption and distribution throughout the body, lithium's half-life may be as high as 18 and 36 hours in the adult and the elderly, respectively⁵.

As explained in the pharmacokinetics section, lithium has several biological interactions that may lead to lithium toxicity. The average patient on chronic lithium treatment may experience at least one episode of toxicity during his lifetime⁶. Lithium toxicity is classified by most authors in three different types: Acute, which includes manifestations that are predominantly gastrointestinal, such as nausea, vomiting and diarrhea, but neuromuscular manifestations may also be present; Acute on chronic, in which both gastrointestinal and neurologic manifestations occur; and Chronic that includes manifestations primarily neurological, including sluggishness, ataxia and neuromuscular excitability⁷.

Toxicity is explained mostly by abnormally reduced excretion in the kidneys, overdose, prescribing errors and drug-to-drug interactions. Lithium has no protein or tissue binding, therefore, hemodialysis provides a useful therapeutic strategy⁸.

At the intracellular level, lithium ion is transported from the lumen to the intracellular space by different protein channels, by means of an exchanging cation pump, in substitution of either sodium or potassium. The H⁺/Na⁺ exchanger accounts for most of the lithium transported into the cytosol and posteriorly to the blood. Particularly, lithium entrance is handled by passive transports compared to its exit, which is handled by pumps involving active transport and energy use. Secondary to this pump imbalance, lithium tends to accumulate throughout the body: Other channels might also contribute to the accumulation of Lithium, such as Glucose/Na channel or Na/K ATPase⁹.

Back to our patient: Our patient developed an urinary tract infection (UTI), particularly a pyelonephritis, as manifested by its systemic symptoms, clinical findings and urinalysis findings. According to classification criteria, our patient met criteria urosepsis etiology. Its urine culture was found to be positive for *Proteus Mirabilis*, a facultative anaerobe bacteria, urease positive with increasing rates of antibiotics resistance¹⁰. Because of the pyelonephritis, induced renal impairment manifested as reduced GFR and an increased reabsorption (secondary to its sepsis induced vasodilation and dehydration), leading to increased lithium levels up to 2.85 mEq/L.

The main stem for treatment of acute lithium toxicity relies on following the ABC protocols (Airway, Breath, Circulation) and supportive care, followed by aggressive hydration in order to guarantee appropriate lithium clearance. Hydration should be done with normal saline (0.9%). Sodium levels should be assessed prior to treatment because these patients may develop induced nephrogenic diabetes insipidus, which makes it important to have a baseline serum sodium level upon hydration. Treatment with oral charcoal is not recommended by most authors¹¹. Some studies have shown a benefit of giving high doses of kayexelate (oral sodium polystyrene sulfonate) is reducing lithium levels, however, its use is not recommended secondary to the side effects of hypokalemia¹². Ultimately, the gold standard treatment when these therapies fail is the use of hemodialysis as an extracorporeal measure. Consensus guidelines for the use of hemodialysis in this setting are controversial. Some authors suggest using hemodialysis in patients with

lithium levels greater than 5 mEq/L and a normal renal function. In those with impaired renal function (creatinine higher than 2 mmol/L) and lithium levels of 4 mEq/L, therapy can be considered as well¹³. Finally, those patients who don't respond to previous treatment and in whom signs of toxicity persist, hemodialysis remains as a reasonable option. More studies are needed to provide guidelines with a stronger degree of recommendation¹⁴.

In our patient, the lithium levels quickly fell after hemodialysis, from 2.85 to 0.89 mEq/L in a 24-hour period. Additionally, the BUN and creatinine levels also fell down to normal after the prerenal acute kidney injury was solved. Creatinine and BUN decreased from 1.89 to 0.85mg/dl and from 43 to 17mg/dl respectively, in a 48-hour period.

Conclusion

Patients treated for bipolar affective disorder with lithium must be specifically warned on how to early identify possible triggers that may induce acute kidney injury, such as UTI, systemic infections, volume depletion states patients must stop medication and seek medical care.

Detailed clinical history of patients initial neurological disorders before initiating Lithium therapy is required, since lithium intoxication symptoms tend to be mostly neurological, so there must be a clear differentiation between the treated pathology symptoms and the newly acquired intoxication symptoms for the diagnosis to be possible and timely. In this case, there were three potential triggers for the GFR drop: Pyelonephritis, prerenal acute kidney injury and prolonged lithium exposure. The summation of these caused Lithium serum levels to increase to toxic levels and resulted in an intoxication, however, the dehydration had the most evident repercussions.

Contribution of each author

Carlos Aníbal Restrepo: In charge of drafting the discussion section compiling the ideas of all the authors. Was also in charge of investigation and evidence to support all the arguments presented.

Maria Antonia Rico: In charge of providing the background information part of the discussion. Also was crucial in the search for evidence and other articles that supported our thesis about the fisiopathology of the intoxication.

Juan Simon Rico: Was the main editor of the paper in charge of retouching and approving the final product. Also contributed with ideas and structure for the development of the whole paper, both discussion and introduction.

Mariana Montoya Castillo, Diana Carolina Echeverri Díaz and Felipe Osorio Ospina: Came up with the idea of presenting the case. Mariana was in charge of collecting all the information regarding our patient. They drafted the introduction of the paper which included all the clinical history of the patient in order to develop a sustained discussion and conclusions.

Conflicts of interest

The authors have no conflicts of interest of any type, nor have any relation with the pharmaceutic industry that could bias the manuscript.

The authors didn't receive financing for the development of the paper and no one else is in any way involved with the article.

Ethical responsibilities

Protection of people and animals

The authors declare that no experiments were performed on human beings or animals for this research.

Data confidentiality

The authors declare that they have followed the protocols of their workplace on the publication of patient data.

The authors certify that we do not have conflicts of interest

No sources of funding were required to carry out the research presented.

The authors confirm that each one contributed with all the material included in the article

References

1. Sturzbecher-Höhne M, Kissner R, Nauser T, Koppenol WH. Preparation and properties of lithium and sodium peroxy nitrite. *Chem Res Toxicol.* 2008;21(12):2257-9. Available from: <https://doi.org/10.1021/tx800279n>
2. Connor H. Drug Discovery—A History. *J R Soc Med.* 2005;98(11):517-8.
3. Jefferson JW, Kalin NH. Serum Lithium Levels and Long-term Diuretic Use. *JAMA.* 1979;241(11):1134.6. Available from: <https://doi.org/10.1001/jama.1979.03290370038025>
4. Gelenberg AJ, Hopkins HS. Report on efficacy of treatments for bipolar disorder. *Psychopharmacol Bull.* 1993;29(4):447-56.
5. Amdisen A. Clinical features and management of lithium poisoning. *Med Toxicol Adverse Drug Exp.* 1988;3(1):18-32.
6. Gitlin M. Lithium side effects and toxicity: prevalence and management strategies. *Int J Bipolar Disord.* 2016;4(1):27. Available from: <https://doi.org/10.1186/s40345-016-0068-y>
7. Offerman SR, Alsop JA, Lee J, Holmes JF. Hospitalized lithium overdose cases reported to the California Poison Control System. *Clin Toxicol.* 2010;48(5):443-8. Available from: <https://doi.org/10.3109/15563650.2010.482533>.
8. Dupuis RE, Cooper AA, Rosamond LJ, Campbell-Bright S. Multiple Delayed Peak Lithium Concentrations following Acute Intoxication with an Extended-Release Product. *Ann Pharmacother.* 1996;30(4):356-60. Available from: <https://doi.org/10.1177/10600280960300406>
9. Timmer RT, Sands JM. Lithium Intoxication. *J Am Soc Nephrol.* 1999;10(3):666-74.
10. Coker C, Poore CA, Li X, Mobley HL. Pathogenesis of *Proteus mirabilis* urinary tract infection. *Microbes Infect.* 2000;2(12):1497-505. Available from: [https://doi.org/10.1016/S1286-4579\(00\)01304-6](https://doi.org/10.1016/S1286-4579(00)01304-6)
11. Linakis JG, Lacouture PG, Eisenberg MS, Maher TJ, Lewander WJ, Driscoll JL, et al. Administration of Activated Charcoal or Sodium Polystyrene Sulfonate (KayexalateTM) as Gastric Decontamination for Lithium Intoxication: An Animal Model. *Pharmacology & Toxicology.* 1989;65(5):387-9. Available from: <https://doi.org/10.1111/j.1600-0773.1989.tb01194.x>
12. Roberge RJ, Martin TG, Schneider SM. Use of sodium polystyrene sulfonate in a lithium overdose. *Ann Emerg Med.* 1993;22(12):1911-5. Available from: [https://doi.org/10.1016/S0196-0644\(05\)80422-6](https://doi.org/10.1016/S0196-0644(05)80422-6)
13. Fenves AZ, Emmett M, White MG. Lithium intoxication associated with acute renal failure. *South Med J.* 1984;77(11):1472-4.
14. Decker BS, Goldfarb DS, Dargan PI, Friesen M, Gosselin S, Hoffman RS, et al. Extracorporeal Treatment for Lithium Poisoning: Systematic Review and Recommendations from the EXTRIP Workgroup. *Clin J Am Soc Nephrol.* 2015;10(5):875-87. Available from: <https://doi.org/10.2215/CJN.10021014>

REVISTA COLOMBIANA DE
NEFROLOGÍA
Volumen 5 No. 1 Enero - Junio de 2018

Contenido / Contents

EDITORIAL

- Infecciones del torrente circulatorio en pacientes en hemodiálisis / *Blood-borne infections in hemodialysis patients*.....15
Rafael Alberto Gómez

ARTÍCULOS ORIGINALES / ORIGINAL INVESTIGATION ARTICLES

- Prevalencia de infección asociada a catéter de hemodiálisis en el Hospital Universitario Clínica San Rafael / *Prevalence of catheter-related haemodialysis infections in Hospital Universitario San Rafael, Bogotá, Colombia*.....17
Javier Gómez, Leonardo Pimienta, Rafael Pino, Maité Hurtado, Mariana Villaveces, Fundación Universitaria Juan N. Corpas, Hospital Universitario Clínica San Rafael, Universidad de La Sabana, Bogotá, Colombia

- Resultados clínicos en receptores de trasplante renal posterior a la conversión a ImTOR / *Clinical outcomes after switch to mTOR inhibitors in kidney transplant recipients*26
Liliana Mesa-Ramírez, Juan C. Gómez-Vega, Jessica Pino-Escobar, Lina M Rivera, Eliana Manzi-Tarapues, Juan G. Posada-Chávez1, Johanna Schweineberg-López, Carlos E. Durán-Rebolledo, Jorge I. Villegas-Otálora, Oscar J. Serrano-Ardila y Luis A. Caicedo-Rusca, Fundación Valle del Lili, Universidad Icesi, Cali Colombia

- Evolución de las glomerulopatías asociadas a la artritis reumatoidea / *Evolution of glomerulopathies associated with rheumatoid arthritis*.....36
José Lucas Daza, Yaroslad De La Cruz, Cintia Marín, Martín Zapata, Fernando Segovia, Luis José Daza, Graciela De Rosa, John Galindo, Universidad de Buenos Aires, Argentina, Universidad Pedagógica y Tecnológica de Colombia, Boyacá, Colombia

ARTÍCULO DE REVISIÓN / REVIEW ARTICLE

- Daño renal asociado a metales pesados: trabajo de revisión / *Renal damage associated with heavy metals: review work*43
Juan Daniel Díaz García, Emmanuel Arceo, Instituto Nacional de Ciencias Médicas y Nutrición “Salvador Zubirán”, Hospital General de México “Eduardo Liceaga”. Ciudad de México, México

- Anomalías congénitas del riñón y del tracto urinario en niños con síndrome de Down / *Congenital anomalies of the kidney and the urinary tract in Down syndrome children*54
Víctor Manuel Mora-Bautista, Universidad Industrial de Santander, Colombia

REPORTE DE CASO / CASE REPORT

Nuevas alternativas para el tratamiento de infección urinaria no complicada, presentación de 2 casos clínicos
/ New alternatives for the treatment of uncomplicated urinary tract infection, presentation of 2 clinical cases

.....61

Andrés F. Reyes, Roberto Ramírez Marmolejo, Universidad Santiago de Cali, Cali, Colombia

Hipertensión renovascular secundaria a hipoplasia congénita de arterias renales en un paciente adulto joven. A propósito de un caso / Renovascular hypertension secondary to congenital hypoplasia of renal arteries in middle adult patient. About a case..... 68
Julián Darío Ñañez Paz, Juan David Orozco Burbano, Natalia Hernández Duque, Hospital Universitario San José, Popayán, Colombia

MANEJO DEL RIESGO DE ENFERMEDAD / MANAGEMENT OF DISEASE RISK

What we do and do not know about women and kidney diseases; Questions unanswered and answers unquestioned: Reflection on World Kidney Day and International Woman's Day. / Salud renal y salud en la mujer: Una oportunidad para mejorar las condiciones de las generaciones actuales y futuras.....74
Giorgia B Piccoli, Mona Alrukaimi, Zhi-Hong Liu, Elena Zakharova, Adeera Levin On behalf of the World Kidney Day Steering Committee, University of Torino, Italy, Centre Hospitalier Le Mans, Le Mans, France, Dubai Medical College, Dubai, United Arab Emirates, Nanjing University School of Medicine, Nanjing, China, Moscow State University of Medicine and Dentistry, Moscow, Russian Federation, University of British Columbia, Vancouver, British Columbia, Canada

Volumen 5 No. 2 Julio - Diciembre de 2018

EDITORIAL

Intoxicación por litio, una verdadera urgencia dialítica /A true dialytic urgency: lithium intoxication.....104
Jorge Eduardo Rico-Fontalvo

ARTÍCULOS ORIGINALES / ORIGINAL INVESTIGATION ARTICLES

Características clínicas e histológicas de las enfermedades parenquimatosas renales en una muestra de biopsias renales obtenidas entre el año 2002 y el 2017 en el departamento de Caldas, Colombia / Clinical and histologic characteristic of renal parenchymal diseases in a renal biopsy sample since 2002 to 2017 in Caldas - Colombia.....107
Héctor Guillermo García Herrera, César Augusto Restrepo Valencia, Carlos Alberto Buitrago Villa, Universidad de Caldas, Caldas, Colombia

Factores asociados a la nefropatía inducida por contraste en pacientes hospitalizados a quienes se les realizó tomografía axial computarizada / Factors associated with contrast-induced nephropathy in hospitalized patients who underwent computed tomography118

Maickol Jhoel Tirado Castro, Liliana Rocío Cataño Vargas, Franklyn Edwin Prieto Alvarado, Universidad Sur-colombiana, Neiva, Colombia

Experiencia en hemodiálisis domiciliaria en España / Spanish home hemodialysis experience.....127

Alejandro Pérez Alba, Javier Reque Santiváñez, Milagros Vázquez Gómez, Ramón Pons Prades, Servicio de Nefrología, Hospital Universitario General de Castellón, Castellón de la Plana, España

Tunneled catheters in femoral vein: does the length makes any difference? / ¿Catéteres femorales tunelizados para hemodiálisis, su longitud hace diferencia?137

César Augusto Restrepo Valencia, José Vicente Aguirre Arango, Carlos Alberto Buitrago Villa, University of Caldas, Colombia

Análisis de pacientes en diálisis peritoneal: factores clínico-epidemiológicos y tipo de transporte peritoneal con recambio hipertónico / Analysis of patients in peritoneal dialysis: clinical-epidemiologic factors and peritoneal transport type with hypertonic replacement146

Fernando Arturo Fajardo-Leitzelar, Manuel Sierra, Daniel Martín Barahona-López, Luis Enrique Sánchez-Sierra, Carlos Felipe Matute-Martínez, Daniel Eduardo Mendoza-Sabillón, Andrea Carolina Núñez-Hernández, Osiris Mariel Serrano-Puerto, Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras

Caracterización de la disfunción tiroidea en adultos con enfermedad renal crónica en diálisis / Characterization of thyroid dysfunction in adults with chronic renal disease in dialysis156

Wilson Fernando Chávez Gómez, Amaury Miguel Ariza García, Enrique Carlos Ramos Clason, Universidad del Sinú, Cartagena, Colombia

Caracterización de la enfermedad renal crónica en adultos mayores / Characterization of chronic renal disease in elderly adults.....166

Julio César Candelaria-Brito, Carlos Gutiérrez-Gutiérrez, Héctor Demetrio Bayarre-Vea, Claudia Acosta -Cruz, Delia Margarita Montes de Oca, Orestes Labrador-Mazón, Policlínico Docente “5 de Septiembre”, Centro de Investigaciones Médicas Quirúrgicas (CIMED), Escuela Nacional de Salud Pública, La Habana, Cuba

Valor de corte del cociente proteinuria/creatininuria predictor de proteinuria = 150 mg/24 h en una muestra de estudiantes argentinos. Utilidad de su aplicación para categorización de la proteinuria / Cut-off value of proteinuria/creatininuria ratio predictor of proteinuria= 150 mg/24h in a sample of argentinean students. Its utility in proteinuria categorization.....179

Cecilia Brissón, Verónica Cuestas, Priscila Prono-Minella, Susana Denner, Verónica Fernández, Rosina Bonifacino-Belzarena, Silvia Marsili, María Eugenia Brissón, Universidad Nacional del Litoral, Santa Fe, Argentina

REPORTE DE CASO / CASE REPORT

A true dialytic urgency: lithium intoxication / Intoxicación por litio, una verdadera urgencia dialítica190

Mariana Montoya-Castillo, Tomás Giraldo-Hinestrosa, Felipe Osorio Ospina, Diana Carolina Echeverri Diaz, Maria Antonia Rico-Mesa, Carlos Aníbal Restrepo, Juan Simon Rico-Mesa, CES University, Pontifical Bolivarian University of Colombia, Cooperative University of Colombia, Medellín, Colombia, Mayo Clinic, Rochester, Cardiology department, United States of America